arXiv:1606.07224v1 [math.GT] 23 Jun 2016

REDUCTION OF BRIDGE POSITIONS ALONG A BRIDGE DISK

JUNG HOON LEE

Abstract. Suppose a knot in a 3-manifold is in n-bridge position. We consider a reduction of the knot along a bridge disk D and show that the result is an $(n-1)$-bridge position if and only if there is a bridge disk E such that (D, E) is a cancelling pair. We apply this to an unknot K, in n-bridge position with respect to a bridge sphere S in the 3-sphere, to consider the relationship between a bridge disk D and a disk in the 3-sphere that K bounds. We show that if a reduction of K along D yields an $(n-1)$-bridge position, then K bounds a disk that contains D as a subdisk and intersects S in n arcs.

1. Introduction

Every closed orientable 3-manifold M can be decomposed into two handlebodies V and W with common boundary S and it is called a *Heegaard splitting* of M, denoted by $M = V \cup_S W$.

For a Heegaard splitting $M = V \cup_S W$, suppose that there are disks $D \subset V$ and $E \subset W$ such that $|D \cap E| = 1$. Then the Heegaard splitting is said to be *stabilized*. A stabilized Heegaard splitting can be destabilized. By compressing V along D (or W along E), we get a lower genus Heegaard splitting. Conversely, it is known that if a compression of V along an essential disk D yields a lower genus Heegaard splitting, then there is a disk E in W such that $|D \cap E| = 1$ [1].

Let K be a knot in M. The notion of Heegaard splitting can be extended to the pair (M, K). For $M = V \cup_S W$, suppose that $V \cap K$ and $W \cap K$ are collections of n boundary parallel arcs. The decomposition $(M, K) = (V, V \cap K) \cup_S (W, W \cap K)$ is called a *bridge splitting* of (M, K), and we say that K is in *n-bridge position* with respect to S. Each arc of $V \cap K$ and $W \cap K$ is a *bridge*. A bridge a cobounds a *bridge disk* D with an arc b in S, i.e. $\partial D = a \cup b$ and $a \cap b = \partial a = \partial b$, such that $D \cap K = a$.

Let a be a bridge in, say, V. An isotopy of a to b along a bridge disk D and further, slightly into W will be called a *reduction* in this paper. We consider when a knot is also in bridge position after a reduction. Not all reductions result in bridge positions. See Figure 1 for an example. A bridge splitting is *perturbed* if there are bridge disks $D \subset V$ and $E \subset W$ such that D intersects E at one point of K. We call (D, E) a *cancelling pair*. It is known that for a perturbed bridge splitting, a reduction along D (or E)

results in a bridge position [4, Lemma 3.1]. We show that the converse is also true. This phenomenon is analogous to the case of Heegaard splitting.

Theorem 1.1. *Let K be a knot in n-bridge position in a 3-manifold M. A reduction of K along a bridge disk D yields an $(n-1)$-bridge position if and only if there is a bridge disk E such that (D, E) is a cancelling pair.*



FIGURE 1. A reduction that does not yield a bridge position

We apply Theorem 1.1 to an unknot in S^3. An unknot K in n-bridge position can be contained in a 2-sphere P that intersects the bridge sphere S in a single loop [2], [3]. Hence each of the two disks that K bounds in P intersects S in n arcs. We study when a disk intersecting S in n arcs contains a given bridge disk, and show that the condition on the bridge disk as in Theorem 1.1 guarantees it. Such a disk containing a bridge disk might be useful to study a collection of disjoint bridge disks, hence furthermore to study a collection of disjoint compressing disks. However, we remark that the converse of Theorem 1.2 does not hold (Figure 1).

Theorem 1.2. *Let K be an unknot in n-bridge position in S^3. If a reduction of K along a bridge disk D yields an $(n-1)$-bridge position, then K bounds a disk F such that*

- *F contains D as a subdisk, and*
- *F intersects the bridge sphere in n arcs.*

In Section 2, we give a proof of Theorem 1.1. In Section 3, we prove Theorem 1.2 using a cancelling pair.

2. Proof of Theorem 1.1

Let $(M, K) = (V, V \cap K) \cup_S (W, W \cap K)$ be a bridge splitting and let K be in n-bridge position. One implication that if (D, E) is a cancelling pair, then

a reduction of K along D yields an $(n-1)$-bridge position is well-known, e.g. [4, Lemma 3.1], so we omit the proof here.

Now we go the other direction. Suppose a reduction of K along a bridge disk D yields an $(n-1)$-bridge position. We assume that D is in V without loss of generality. Denote $D \cap K$ by a and $D \cap S$ by b. By a reduction along D, a is isotoped past b to an arc a' in W. Consider the rectangle R with opposite sides a' and b. It is a part in W of the region which a swept out. See Figure 2. We denote the other two sides of R by α and β.



FIGURE 2. A reduction along D

The arc a' is a subarc of a new bridge c. We denote a bridge disk for c by C and let $d = C \cap S$. The bridge disk C and the rectangle R have a' in common and they may have other intersection. Suppose that there exists an arc component of $R \cap C$ with an endpoint in $\operatorname{int} a'$. This happens if, around the axis a', C spirals with respect to R. Let γ be such an arc with a point x of $\partial\gamma$ closest to the point $y = a' \cap \alpha$. We push a collar neighborhood of the line segment $\overline{xy}$ in C to the opposite side of R as in Figure 3. Then γ is changed into an arc with an endpoint in $\operatorname{int} \alpha$. In this way, even if C spirals more than $360°$, we isotope every arc component of $R \cap C$ with an endpoint in $\operatorname{int} a'$ so that the endpoint is not in $\operatorname{int} a'$. Hence we may assume that there exists no arc component of $R \cap C$ with an endpoint in $\operatorname{int} a'$. (In fact, we do not necessarily need to isotope an endpoint into $\operatorname{int} \alpha$ (or $\operatorname{int} \beta$). See Figure 4 for an example.)



FIGURE 3. An isotopy of C



FIGURE 4. Two arcs are merged into one.

A circle component of $R \cap C$ can be removed out of R by an isotopy passing through α or β. An arc component of $R \cap C$ belongs to one of the six types below according to its endpoints.

(1) Both endpoints are in α.
(2) Both endpoints are in β.
(3) Both endpoints are in b.
(4) One endpoint is in α and the other is in b.
(5) One endpoint is in β and the other is in b.
(6) One endpoint is in α and the other is in β.

Every arc of types (1), (2), (3), (4), (5) can be removed out of R by an isotopy (Figure 5), but an arc of type (6) cannot be removed. So we may assume that $R \cap C$ consists of type (6) arcs and the number of components $k = |R \cap C| - 1$ is minimal. See Figure 6. It will be shown that $k = 0$. So for this purpose we suppose that $k > 0$.



FIGURE 5. Components of $R \cap C$ that can be removed

Now we reverse the reduction operation. The arc a' moves back to $\alpha \cup a \cup \beta$ along $R \cup D$ and the bridge disk C moves together. Of course, some neighborhoods of the k arcs of $R \cap C$ are pushed into V, producing k parallel lower caps for D. See Figure 7. After this isotopy, C contains D as a subdisk and $C \cap W$ is a k-punctured disk. Let $C' = C \cap W$.

Let e and f be the two bridges in W that are adjacent to the bridge a in the original n-bridge position of K. We denote a bridge disk for e by E. We choose E such that $|C' \cap E|$ is minimal.



FIGURE 6. The intersection $R \cap C$



FIGURE 7. The disk C after reversing the reduction operation

The component of $\partial C'$ which is not a puncture can be regarded as a 4-gon consisting of two bridges e and f and two arcs b and d in S. We give label 1 to the puncture of C' which is closest to b in S, and label consecutively the other nested punctures in S by $2, \dots, k$. See Figure 7 and Figure 8.

If E spirals with respect to C' around the axis e, then as before we isotope an endpoint of an arc of $C' \cap E$ in $\operatorname{int} e$ into $\operatorname{int} d$. See Figure 9. So we may assume that there is no arc component of $C' \cap E$ with an endpoint in $\operatorname{int} e$.

Remark 2.1. After the isotopy in Figure 9, the arc $E \cap S$ does not intersect d minimally in $S - K$. In other words, there is a bigon formed by $(E \cap S) \cup d$ bounding a disk in $S - K$. But what is needed in our argument is that $(E \cap S) \cup b$ does not make a bigon bounding a disk in $S - K$.

Lemma 2.2. *The intersection $C' \cap E$ consists of arcs, each of which is either*

- *the arc e, or*
- *not boundary parallel in C', or*
- *boundary parallel in C' with one endpoint in b and the other in d.*



FIGURE 8. The k-punctured disk C'



FIGURE 9. An isotopy of E

Proof. Suppose that there is a circle component of $C' \cap E$ which is inessential in C'. (Here, inessential means that it bounds a disk.) Let γ be an innermost such component in C' and let Δ be the corresponding innermost disk. Then a disk surgery of E along Δ reduces $|C' \cap E|$, a contradiction. Hence all circle components of $C' \cap E$ are essential in C'. Let γ be an innermost circle component of $C' \cap E$ in E and Δ be the corresponding innermost disk. By a disk surgery of C' along Δ, we get a disk that replaces C' and has fewer punctures, which contradicts the minimality of the number of punctures. So we may assume that $C' \cap E$ consists of arc components.

Suppose that there exists an arc of $C' \cap E$ (except for e) such that the arc is boundary parallel in C' and it is not the case that one endpoint of the arc is in b and the other endpoint is in d. Then both endpoints are in b, or in d, or in a puncture. We choose an outermost such arc γ in C' and let Δ be the corresponding outermost disk. A surgery of E along Δ reduces $|C' \cap E|$, a contradiction. So we conclude that any boundary parallel arc of $C' \cap E$ in C' has one endpoint in b and the other in d. □

We give each endpoint of arcs of $C' \cap E$ a label among $\{b, d, 1, 2, \ldots, k\}$. Consider a sequence of labels that appears along $E \cap S$. It begins with b and $1, 2, \ldots, k$ follow by Remark 2.1 and because the k punctures in S are nested around b. Then a possibly empty subsequence of d's follows and then $k, \ldots, 1$ follow and then b follows again. This pattern is repeated and the sequence ends with d. See Figure 10 for an appearance of the sequence. In the figure, (d) means that it can be empty. We consider a pair of labels of an arc of $C' \cap E$ without order.



FIGURE 10. A sequence of labels on $E \cap S$

Lemma 2.3. *A pair of labels of an outermost arc of $C' \cap E$ in E is (k, k).*

Proof. Let γ be an outermost arc of $C' \cap E$ in E and let Δ be the corresponding outermost disk. By the above paragraph, the pair of labels of $\partial\gamma$ is one of the following.

(1) $(b, 1)$
(2) $(i, i+1)$ $(i = 1, \ldots, k-1)$
(3) (k, d)
(4) (d, d)
(5) (k, k) (There is no d between consecutive two k's.)

Case (1) $(b, 1)$
By a boundary compression of C' along Δ, the arc b and the puncture with label 1 is connected into an arc, which is still isotopic to b in $S - K$. In V, we can see that D and the lower cap disk is connected by a band, yielding a disk isotopic to D. See Figure 11. Hence there exists a disk with fewer punctures replacing C', a contradiction.

Case (2) $(i, i+1)$ $(i = 1, \ldots, k-1)$
In this case, a boundary compression of C' along Δ changes puncture i and $i+1$ into an inessential loop in $S - K$. So we can reduce the number of punctures by two, a contradiction.

Case (3) (k, d)
A boundary compression of C' along Δ connects d and puncture k into an arc. This also reduces the number of punctures, a contradiction.



FIGURE 11. A band sum of D and the lower cap disk is isotopic to D.

Case (4) (d, d)
Since γ is not a boundary parallel arc in C', a boundary compression of C' along Δ reduces the number of punctures, a contradiction.

So the pair of labels of $\partial\gamma$ should be the remaining Case (5). An appearance of the sequence of labels is as in Figure 12. □



FIGURE 12. A sequence of labels on $E \cap S$

Lemma 2.4. $k = 0$.

Proof. Consider only all outermost arcs of $C' \cap E$ in E. It is easy to see that there are more than one outermost arcs. We choose two outermost arcs α and β whose endpoints are consecutive along $E \cap S$ such that every other arc with one of its endpoints between $\partial\alpha$ and $\partial\beta$ are parallel to α or β as in Figure 13. By the pattern of the sequence of labels, all labels $1, \ldots, k-1$ appear between $\partial\alpha$ and $\partial\beta$. So an arc γ_i with a pair of boundary labels (i, i) exists for all i $(i = 1, \ldots, k)$. If we cap off each puncture i of C' with a disk Δ_i, we get a disk $\overline{C}$ from C'. Then $\Delta_i \cup \gamma_i$ can be regarded as a loop with fat vertex in $\overline{C}$. Consider an innermost one $\Delta_j \cup \gamma_j$ in $\overline{C}$. Then γ_j cuts off a

disk from C' and this contradicts Lemma 2.2 that an arc with label (i, i) is not bounday parallel in C'. The contradiction is caused by the assumption that $k > 0$. So we conclude that $k = 0$ and C' is a disk. □



FIGURE 13. Two outermost arcs α and β in E

By Lemma 2.4, C' is a disk. Again we choose E such that $|C' \cap E|$ is minimal. Suppose that $|C' \cap E| > 0$. If E spirals with respect to C', then we isotope an endpoint of an arc of $C' \cap E$ in $\operatorname{int} e$ into $\operatorname{int} b$ or $\operatorname{int} d$. By similar argument as in the proof of Lemma 2.2, $C' \cap E$ (except for e) consists of arcs, each with one endpoint in b and the other in d. See Figure 14. Let γ be an outermost arc of $C' \cap E$ in C' cutting off an outermost disk Δ containing e. Let E_1 and E_2 be the two subdisks of E cut off by γ with $e \subset E_1$. Then $\Delta \cup E_2$ is a new bridge disk for e with $|C' \cap (\Delta \cup E_2)| < |C' \cap E|$, a contradiction. Therefore $C' \cap E = e$ and this implies that $D \cap E$ is a single point. This completes the proof of Theorem 1.1.



FIGURE 14. The arcs of $C' \cap E$ in C'

3. Proof of Theorem 1.2

By Theorem 1.1, there exists a bridge disk E such that (D, E) is a cancelling pair. Let $a = D \cap K$ and $b = E \cap K$ and $p = D \cap E$. We simultaneously isotope a and b along D and E respectively and further slightly into W and

V respectively, fixing the point p. Let a' be the resulting arc of a after the isotopy and let Δ_1 be the region that a swept out. Then Δ_1 contains D. Similarly, let b' be the resulting arc of b after the isotopy and let Δ_2 be the region that b swept out.

The unknot K is in $(n-1)$-bridge position now and bounds a disk F_0 that intersects S in $n-1$ arcs. The disk F_0 and $\Delta_1 \cup \Delta_2$ have a' and b' in common and they may have other intersection. If F_0 spirals with respect to Δ_1 (or Δ_2) around the axis a' (or b') respectively, then we isotope F_0 as before so that there is no arc component of $(\Delta_1 \cup \Delta_2) \cap F_0$ with an endpoint in $\operatorname{int} a'$ or $\operatorname{int} b'$. We move all circle components and arc components (except for a' and b') of $(\Delta_1 \cup \Delta_2) \cap F_0$ horizontally to the outside of $\Delta_1 \cup \Delta_2$ so that the property of $F_0 \cap S$ being $n-1$ arcs is preserved. See Figure 15.



FIGURE 15. Horizontal isotopy of F_0

Now we give a perturbation to K using Δ_1 and Δ_2. The boundary of $F = F_0 \cup \Delta_1 \cup \Delta_2$ is the original K in n-bridge position. But $F \cap S$ is not a collection of n arcs. We slightly push a part of F in V into W as in Figure 16 so that $F \cap S$ is a collection of n arcs and F still contains D as a subdisk.

Acknowledgements. The author would like to thank Toshio Saito for helpful discussion.

REFERENCES

[1] **C. McA. Gordon**, *On primitive sets of loops in the boundary of a handlebody*, Topology Appl. **27** (1987) 285–299.
[2] **J. -P. Otal**, *Présentations en ponts du nœud trivial*, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982) 553–556.
[3] **M. Ozawa**, *Bridge position and the representativity of spatial graphs*, Topology Appl. **159** (2012) 936–947.
[4] **M. Scharlemann and M. Tomova**, *Uniqueness of bridge surfaces for* 2*-bridge knots*, Math. Proc. Cambridge Philos. Soc. **144** (2008) 639–650.

DEPARTMENT OF MATHEMATICS AND INSTITUTE OF PURE AND APPLIED MATHEMATICS, CHONBUK NATIONAL UNIVERSITY, JEONJU 54896, KOREA
E-mail address: junghoon@jbnu.ac.kr



FIGURE 16. An isotopy of F